

Toward Mapping the Detailed Density Structure of Classical Be Circumstellar Disks

J.P. Wisniewski^{1,2,3}, A.F. Kowalski^{4,5}, K.S. Bjorkman^{3,6}, J.E. Bjorkman^{3,6}, & A.C. Carciofi⁷

ABSTRACT

The first results from a near-contemporaneous optical and infrared spectroscopic observing program designed to probe the detailed density structure of classical Be circumstellar disks are presented. We report the discovery of asymmetrical infrared emission lines of He I, O I, Fe II, and the Brackett, Paschen, and Pfund series lines of H I which exhibit an opposite V/R orientation ($V > R$) to that observed for the optical Balmer H α line ($V < R$) in the classical Be star ζ Tau. We interpret these data as evidence that the density wave which characterizes ζ Tau's disk has a significantly different average azimuthal morphology in the inner disk region as compared to the outer disk region. A follow-up multi-wavelength observational campaign to trace the temporal evolution of these line profile morphologies, along with detailed theoretical modeling, is suggested to test this hypothesis.

Subject headings: stars: emission-line, Be — circumstellar matter — stars: individual (ζ Tau) — stars: individual (48 Librae)

¹NASA Goddard Space Flight Center Code 667 Greenbelt, MD 20771, jwisnie@milkyway.gsfc.nasa.gov

²NPP Fellow

³Visiting Astronomer at the Infrared Telescope Facility, which is operated by the University of Hawaii under Cooperative Agreement no. NCC 5-538 with the National Aeronautics and Space Administration, Science Mission Directorate, Planetary Astronomy Program

⁴Department of Physics, University of Chicago 5640 S. Ellis Ave. Chicago, IL 60637

⁵Current address: Department of Astronomy, University of Washington Box 351580 Seattle, WA 98195, kowalski@astro.washington.edu

⁶Department of Physics & Astronomy, University of Toledo MS113, Toledo, OH 43606 USA, karen@astro.utoledo.edu, jon@physics.utoledo.edu

⁷Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, Rua do Matão 1226, Cidade Universitária, São Paulo, SP 05508-900, Brazil, carciofi@usp.br

1. Introduction

Both observational and theoretical evidence indicates that classical Be stars are near main sequence B-type stars with gaseous circumstellar disks (e.g. see review by Porter & Rivinius 2003); however, the mechanism(s) responsible for creating these disks are not well understood. Optical spectroscopic studies have uncovered a myriad of different line profile morphologies commonly observed in classical Be stars (Hanuschik et al. 1996), including double-peaked asymmetrical emission lines (McLaughlin 1961; Dachs et al. 1992; Hanuschik et al. 1995). The ratio of the violet (V) to red (R) flux within individual lines, hereafter referred to as a V/R ratio, is observed to vary on quasi-periodic timescales of several to ten years, as documented for a large set of Be stars by Okazaki (1997). While numerous theories have been proposed to explain this observed phenomenon, as summarized within Hanuschik et al. (1995), these features are most commonly interpreted as evidence of one-armed density waves within these circumstellar disks (Okazaki 1991; Hanuschik et al. 1995; Okazaki 1997). Note however that the presence of phase lags between polarimetric and $H\alpha$ observations has led McDavid et al. (2000) to suggest these waves might be more spiral-like than previously thought. Interferometric studies (Vakili et al. 1998; Berio et al. 1999) have also provided evidence that Be disks deviate from axisymmetry, namely they have provided evidence of prograde one-armed oscillations in several disks.

It is not uncommon to observe multiple emission lines in a spectrum of a classical Be star that exhibit a V/R asymmetry, and with few exceptions the same orientation of this asymmetry is observed in all lines. Baade (1985) summarizes several of these exceptions, in which a phase lag appears to be present between optical hydrogen and Fe II lines (Slettebak 1982) and between different lines in the hydrogen Balmer series (Kogure & Suzuki 1984). Baade (1985) also suggested that a phase lag exists between the $H\alpha$ and hydrogen Pa 15-18 lines of γ Cas presented by Chalabaev & Maillard (1983); however, we believe this suggestion is rather dubious given the poor resolution of these data and clear evidence of contamination from nearby lines.

Few moderate resolution infrared (IR) spectroscopic studies of classical Be stars have been performed to date. Clark & Steele (2000) and Steele & Clark (2001) have investigated a large set of Be stars in the H and K band at sufficient resolution to begin to search for detailed line profile effects. The first tentative evidence of IR V/R reversals, analogous to the optical V/R phase lags summarized by Baade (1985), was presented in Clark & Steele (2000), who noted that the Brackett γ and He I $2.058\mu\text{m}$ lines had opposite orientations in one of their targets.

In this Letter, we present the initial results from a near-contemporaneous, optical and IR spectroscopic observing campaign designed to provide insight into the detailed density

structure of classical Be circumstellar disks, hence diagnose the mechanisms responsible for creating these disks. Detailed modeling of these data, as well as follow-up observations, will be presented in subsequent papers. In Section 2 we outline the observational data to be discussed in this work. An analysis and discussion of these data are presented in Section 3. Finally, a summary of our main results and suggested follow-up work is presented in Section 4.

2. Observations

The infrared (IR) spectroscopic data presented in this paper were obtained at the 3m NASA Infrared Telescope Facility (IRTF), using the SpeX medium resolution spectrograph (Rayner et al. 2003). We used a 0.3×15 arc-second slit to provide wavelength coverage from $0.8\text{--}2.4\mu\text{m}$ at $R \sim 2000$. Observations of A0V stars were obtained at ~ 15 minute intervals throughout our run, at a similar hour angle and airmass to our science targets, to serve as telluric standards. Further details regarding the telluric correction method commonly applied to SpeX data are presented in Vacca et al. (2003). All observations were reduced using Spextool (Cushing et al. 2004). These data were taken during a period of extremely poor weather conditions on Mauna Kea, thus we have not attempted to perform a rigorous flux calibration. While these observing conditions did introduce small artifacts in wavelength regimes outside of those presented here (see Wisniewski 2005), we are fully confident that the line profile morphologies and line-to-continuum ratios of the data presented in this paper are not influenced by the presence of any residual telluric features.

The optical spectroscopic data presented here were obtained at the 1m telescope of the Ritter Observatory, using its fiber-fed echelle spectrograph. Nine non-overlapping orders, each of width $70 \text{ \AA}$, were obtained over the wavelength range $5300\text{--}6600\text{\AA}$ at $R \sim 26000$. Standard IRAF¹ techniques were used to reduce these data, including procedures to apply bias, flat field, and wavelength calibration corrections to all data. Further details regarding the reduction of Ritter data can be found in Morrison et al. (1997). A summary of the optical and IR data to be discussed in this paper are presented in Table 1. Note that we have not corrected any of our optical or IR spectral line data for underlying photospheric absorption components.

¹IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under contract with the National Science Foundation.

3. Results and Discussion

As demonstrated by our near-contemporaneous optical and IR observations of 48 Librae (see Figure 1), double-peaked asymmetrical profiles in Be spectra often exhibit the same V/R orientation for all observed transitions. Slettebak (1982) reported phase lags between optical Fe II and hydrogen lines in several of his observations of Be stars; we found evidence of a similar phenomenon in our IR observation of the suggested classical Be star NGC 2439:WBBel (see Wisniewski 2005; Wisniewski & Bjorkman 2006).

We obtained two H α spectra of the well studied classical Be star ζ Tau (see Figure 2) one month before and one month after our IR observations, and note that both H α profiles exhibit triple-peaked profiles characterized by an overall V < R asymmetry. The V/R period is often several years, so the line profiles are generally stable on timescales of several months (Okazaki 1997). We are confident that ζ Tau exhibited a H α profile similar to that shown in Figure 2 at the time of our IR observation, and was not in the midst of a cycle of irregular fluctuations of the type described by Guo et al. (1995), based on our long baseline of previous observations of the star. At the resolution of SpeX, ζ Tau’s IR spectrum shows many emission-lines characterized by a range of morphologies, including single-peaked profiles, asymmetrical double-peaked profiles, and asymmetrical double-peaked shell profiles. As shown in Figure 2, hydrogen Brackett transitions in these data exhibit shell-like profiles, which are asymmetrical with V > R. Other IR emission-lines which have asymmetrical profiles with a similar V/R orientation to the H I Brackett series include O I 0.8446, 1.129, 1.317 μm ; Fe II 1.086, 1.113 μm ; He I 1.083 μm ; H I Paschen γ and β ; and the H I Pfund lines at 2.382, 2.392, 2.403 μm . These data exhibit a fundamentally different V/R morphology than our near-contemporaneous H α profile.

As ζ Tau is known to experience cyclical V/R variations, likely due to the presence of a density wave in its disk (Okazaki 1997), we suggest that the observed opposite V/R orientation of the IR lines and H α can be explained by the presence of a density wave in the disk which has a significantly different average azimuthal morphology (e.g. a spiral wave), combined with different loci for line formation. This idea is illustrated in Figure 3, where we show schematically the disk of ζ Tau with a one-armed spiral density perturbation: note that lighter shading denotes regions of density enhancement while darker shading denotes density decrements. The H I IR lines are much more optically thin than the optical lines, hence they form within the inner disk, while the optical H I lines originate from a much larger area. As a result of the different azimuthal properties of the disk with respect to the distance from the star, in certain phases of the V/R cycle the net density enhancement of the inner disk may occur in an opposite side (with respect to the line of sight) to the net density enhancement of the entire disk, thus producing line profiles with opposite V/R ratios. For instance, in

Figure 3 the $V > R$ morphology of Br-12 is produced by a net density enhancement in the inner disk region which is moving towards the observer, and the $V < R$ morphology of $H\alpha$ is produced by a net density enhancement in the outer disk region which is moving away from the observer.

Carciofi & Bjorkman (2006), Carciofi et al. (2006), and Carciofi et al. (2007) outline a new Monte Carlo code for solving the radiative transfer in the circumstellar disks of Be stars. This fully three-dimensional non-LTE Monte Carlo code produces a self-consistent determination of the hydrogen level populations, electron temperature, and gas density of a circumstellar disk, given a prescription of the central star and assuming a static, nearly-Keplerian disk with a given mass loss rate, truncation radius, and a prescription for the gas viscosity. In Carciofi et al. (2005), the authors used this code to study the circumstellar disk of ζ Tau. Following Wood et al. (1997), they adopted a luminosity class of IV and a spectral type of $B2.9 \pm 0.4$, which corresponds to an effective temperature of 19,000K and a stellar radius of $5.6R_{\odot}$. Assuming an outer disk radius of $100 R_{\star}$ and a mass loss rate of $2.5 \times 10^{-11} M_{\odot}/\text{yr}$ the authors were able to fit the observed SED from visible to mid IR wavelengths, and optical ($\lambda\lambda$ 3200–10500 Å) spectropolarimetry. Using this model, we calculated intensity maps of $H\alpha$ and Br-12 for ζ Tau which, as seen in Figure 4, illustrate that Br-12 is formed in the inner disk ($R \lesssim 15R_{\star}$) whereas $H\alpha$ is produced over a much more extended region, thereby confirming our previous assertion that our optical versus IR H I data probe distinctly different disk regions. Future iterations of the model will incorporate complex density profiles, such as one-armed waves, which will enable us to test our exploratory discussion in a more quantitative manner.

We note that V/R reversals have been predicted by at least one other theory in the refereed literature. Waters & Marlborough (1992) predicted instances in which a ratio of $R > V$ in a $H\alpha$ line would reverse to a ratio of $R < V$ in IR H I lines, due to line self-absorption in the inner region of edge-on disks. However, the presence of this reversal depends critically on the existence of a large expansion component in the disk velocity at the location where the lines form, an unlikely scenario given the very small radial velocities now believed to characterize inner disk regions (Rivinius et al. 1999; Hanuschik 2000; Porter & Rivinius 2003).

4. Summary and Future Work

We have presented the first results of near-contemporaneous optical and IR spectroscopic observations of classical Be stars. We report the discovery of asymmetrical infrared emission lines (He I, O I, Fe II, H I Paschen, Brackett, and Pfund series) in ζ Tau which have oppositely

oriented V/R ratios to that seen at $H\alpha$. We suggest these data indicate that the density wave which characterizes ζ Tau’s disk has a significantly different average azimuthal morphology in the inner disk region as compared to its outer disk region. This would also indicate that the Fe II lines are formed in the inner disk, in the same spatial location as the IR lines are formed. These features should travel through the disk on the cyclical timescale generally reported for density waves in ζ Tau (Okazaki 1997); hence, we would predict continued anti-correlation amongst these line profiles over the course of several observing seasons. We have begun a long-term optical and IR spectroscopic monitoring program to trace this evolution. As demonstrated by the initial results presented in this Letter, such an observational approach has the potential to provide powerful constraints to theoretical efforts to map the detailed density distribution of Be circumstellar disks, hence better understand the mechanism(s) which create these disks.

We thank our referee, Myron Smith, for comments which helped to improve this manuscript. We also thank the staff of Ritter Observatory for their assistance in providing data presented in this paper. This research was supported by a NASA LTSA grant to KSB (NAG5-8054), a Research Corporation Cottrell Scholar award (KSB), a NASA GSRP Fellowship to JPW (NGT5-50469), a Sigma Xi grant to JPW, and the NSF REU program at the University of Toledo (AFK). JEB and ACC acknowledge support from NSF grant AST-0307686; ACC is also supported by FAPESP grants 01/12589-1 and 04/07707-3. Observations at the Ritter Observatory are supported by the NSF under the PREST program, grant number AST-0440784.

REFERENCES

- Baade, D. 1985, *A&A*, 148, 59
- Berio, P. 1999, *A&A*, 345, 203
- Bjorkman, J.E. & Carciofi, A.C. 2005, in *Astronomical Polarimetry: Current Status and Future Directions*, ed. A. Adamson, C. Aspin, C. J. Davis, & T. Fujiyoshi, ASP Conf. Ser. 343, 417
- Carciofi, A.C. & Bjorkman, J.E. 2006, *ApJ*, 639, 1081
- Carciofi, A.C. et al. 2006, *ApJ*, 652, 1617
- Carciofi, A.C., Bjorkman, J.E., Miroshnichenko, A.S., Magalhães, A.M., & Bjorkman, K.S.

- 2007, in *Active OB Stars: Laboratories for Stellar and Circumstellar Physics*, eds. S. Owocki, S. Stefl & A. Okazaki (ASP Conf. Series 361; San Francisco: ASP), in press
- Chalabaev, A. & Maillard, J.P. 1983, *A&A*, 127, 279
- Clark, J.S. & Steele, I.A. 2000, *A&AS*, 141, 65
- Cushing, M.C., Vacca, W.D., & Rayner, J.T. 2004, *PASP*, 116, 362
- Dachs, J., Hummel, W., & Hanuschik, R.W. 1992, *A&AS*, 95, 437
- Guo, Y., Huang, L., Hao, J., Cao, H., Guo, Z., & Guo, X. 1995, *A&AS*, 112, 201
- Hanuschik, R.W., Hummel, W., Dietle, O., & Sutorius, E. 1995, *A&A*, 300, 163
- Hanuschik, R.W., Hummel, W., Sutorius, E., Dietle, O., & Thimm, G. 1996, *A&AS*, 116, 309
- Hanuschik, R.W. 2000, in *IAU Colloq. 175, The Be Phenomenon in Early-Type Stars*, ed. M.A. Smith, H.F. Henrichs, & J. Fabregat (ASP Conf. Serl. 214; San Francisco: ASP), 518
- Kogure, T. & Suzuki, M. 1984, *PASJ*, 36, 191
- McDavid, D., Bjorkman, K.S., Bjorkman, J.E., & Okazaki, A.T. 2000, in *IAU Colloq. 175, The Be Phenomenon in Early-Type Stars*, ed. M.A. Smith, H.F. Henrichs, & J. Fabregat (ASP Conf. Serl. 214; San Francisco: ASP), 460
- McLaughlin, D.B. 1961, *JRASC*, 55, 73
- Morrison, N.D., Knauth, D.C., Mulliss, C.L., & Lee, W. 1997, *PASP*, 109, 676
- Okazaki, A.T. 1991, *PASJ*, 43, 75
- Okazaki, A.T. 1997, *A&A*, 318, 548
- Porter, J.M. & Rivinius, T. 2003, *PASP*, 115, 1153
- Quirrenbach, A. et al. 1997, *ApJ*, 479, 477
- Rayner, J.T. et al. 2003, *PASP*, 115, 362
- Reig, P., Negueruela, I., Coe, M.J., Fabregat, J., Tarasov, A.E., & Zamanov, R.K. 2000, *MNRAS*, 317, 205

- Rivinius, Th., Stefl, S., & Baade, D. 1999, *A&A*, 348, 831
- Slettebak, A. 1982, *ApJS*, 50, 55
- Steele, I.A. & Clark, J.S. 2001, *A&A*, 371, 643
- Telting, J.H., Waters, L.B.F.M., Persi, P., & Dunlop, S.R. 1993, *A&A*, 270, 355
- Vacca, W.D., Cushing, M.C., & Rayner, J.T. 2003, *PASP*, 115, 389
- Vakili, F. et al. *A&A*, 335, 261
- Waters, L.B.F.M. & Marlborough, J.M. 1992, *A&A*, 253, L25
- Wisniewski, J.P. 2005, Ph.D. Thesis, University of Toledo
- Wisniewski, J.P. & Bjorkman, K.S. 2006, *ApJ*, 652, 458
- Wood, K., Bjorkman, K.S., & Bjorkman, J.E. 1997, *ApJ*, 477, 926




Fig. 1.— *Left:* The H α spectrum of 48 Librae observed on 2004 March 11 exhibits an asymmetrical profile characterized by $V > R$. *Right:* A near contemporaneous infrared spectrum obtained on 2004 March 6 also exhibits hydrogen Brackett lines having asymmetrical profiles with $V > R$. The agreement between the orientation of these line profile morphologies is the standard scenario expected in observations of classical Be stars.




Fig. 2.— *Left:* The H α spectrum of ζ Tau obtained on 2004 February 28 (left panel) clearly exhibits an asymmetrical triple-peaked profile characterized by $V < R$. Although not shown here, an optical spectrum obtained on 2004 April 12 exhibits the same morphology. *Right:* This IR spectrum of ζ Tau obtained on 2004 March 5 is characterized by asymmetrical hydrogen Brackett lines having $V > R$. Although not picture here, additional He I, O I, Fe II, and H I (Paschen and Pfund series) emission lines in this spectrum exhibit this same $V > R$ morphology.




Fig. 3.— A one-armed density perturbation, characterized by regions of density enhancement (lighter regions) and regions of density decrements (darker regions), is believed to reside in ζ Tau’s disk based on observed V/R line profile variations. Our IR (left panel) and optical (right panel) spectroscopic data probe different physical locations within the ζ Tau disk, hence probe different regions of the density perturbation located within the disk.




Fig. 4.— Carciofi & Bjorkman (2006) model images of the ζ Tau disk at H α (upper panel) and H I Br-12 (lower panel). The scale of these images is 40 stellar radii (~ 224 solar radii). These images clearly demonstrate that H α is produced over a much larger disk area, whereas H I Br-12 is produced from a more inner disk region.

Table 1: Summary of Observations

Object	Observatory	Date	Exposure Times
ζ Tau	I	2004 March 5	60 s.
ζ Tau	R	2004 February 28	1200 s.
ζ Tau	R	2004 April 12	1200 s.
48 Lib	I	2004 March 6	150 s.
48 Lib	R	2004 March 11	3600 s.

Note. — Some properties of our observational data are summarized. The observatory abbreviation “I” corresponds to IRTF data and “R” corresponds to Ritter data. For our infrared observations, the listed exposure times are the total effective integration times of the science targets, and do not represent the individual exposure times at each of the telescope nod positions described in the text.